



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 31/00	A2	(11) International Publication Number: WO 99/48482 (43) International Publication Date: 30 September 1999 (30.09.99)
(21) International Application Number: PCT/GB99/00960 (22) International Filing Date: 26 March 1999 (26.03.99) (30) Priority Data: 9806513.9 26 March 1998 (26.03.98) GB 9905275.5 8 March 1999 (08.03.99) GB (71) Applicant (for all designated States except US): PHY- TOPHARM PLC [GB/GB]; Corpus Christi House, 9 West Street, Godmanchester, Cambridgeshire PE18 8HG (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): XIA, Zongqin [CN/CN]; Flat 403, Block 8, 1 Jian-De Road, Shanghai 200025 (CN). HU, Yaer [CN/CN]; Flat 1209, Block 1, Lane 357, Ma-Dang Road, Sanghai 200025 (CN). RUBIN, Ian [GB/GB]; Hall Farm House, 9 High Street, Castle Donington, Leicester LE74 2PP (GB). BROSTOFF, Jonathan [GB/GB]; 34 Fitzjohn Avenue, London NW3 5NB (GB). WHITTLE, Brian [GB/GB]; Mereclose, Hull Road, Hornsea, East Yorkshire HU18 1RJ (GB). WANG, Weijun [CN/GB]; 10 Brecon Way, Hinchingsbrooke Park, Huntingdon, Cambs PE18 8XX (GB). GUNNING, Phil [GB/GB]; 37a King Street, Saffron Walden, Essex CB10 1EU (GB).	(74) Agent: ABRAMS, Michael, John; Haseltine Lake & Co., Imperial House, 15-19 Kingsway, London WC2B 6UD (GB). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	
(54) Title: SMILAGENIN AND ANZUROGENIN-D AND THEIR USE		
(57) Abstract		
The invention discloses the use of a smilagenin and anzurogenin-D in the treatment of cognitive disfunction and similar conditions. Methods of treatment, and pharmaceutical compositions are also disclosed.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SMILAGENIN AND ANZUROGENIN-D AND THEIR USE

The present invention relates to smilagenin and anzurogenin-D and their use in treating cognitive dysfunction and allied conditions; and to compositions for use in such treatments. The invention is also concerned with the treatment of conditions that are characterised by a deficiency in the number or function of membrane-bound receptors. In the following, the present invention will be described principally with reference to the treatment of Alzheimer's disease (AD) and senile dementia of the Alzheimer's type (SDAT), where deficiencies in a number of receptor types have been demonstrated. However, it is to be understood that the present invention relates generally to the treatment of conditions attributable to intrinsic pathological conditions and/or exposure to adverse environmental conditions these conditions being characterised by a deficiency in the number or function of membrane-bound receptors or a deficiency in transmission at the junctions between neurones or at the junctions of neurones and effector cells.

Conditions of the type mentioned above include Parkinson's disease, Lewi body dementia, postural hypotension, autism, chronic fatigue syndrome, Myasthenia Gravis, Lambert Eaton disease, diseases and problems associated with Gulf War Syndrome, occupational exposure to organophosphorus compounds and problems associated with ageing.

Alzheimer's disease (AD) and senile dementia of the Alzheimer's type (SDAT) are grave and growing problems in all societies where, because of an increase in life expectancy and control of adventitious disease, the demographic profile is increasingly extending towards a more aged population. Agents which can treat, or help in the management of, AD/SDAT are urgently required.

Age-associated memory impairment (AAMI) is a characteristic of older patients who, while being psychologically and physically normal, complain of

memory loss. It is a poorly defined syndrome, but agents which are effective in treatment of AD/SDAT may also be of value in these patients.

Research into AD/SDAT is being carried out by traditional and
5 conventional medical research methods and disciplines. In conventional
medicine, there are several approaches to the treatment of AD/SDAT. It is
known that the biochemical processes subserving memory in the cerebral cortex
are (at least in part) cholinergically-mediated. Those skilled in the art will
know that "cholinergically mediated" mechanisms may be directly attributable to
10 acetylcholine acting on receptors, and these are direct effects. Other, clinically
useful effects may also be caused by modulation of release of acetylcholine
from pre-synaptic nerve endings or inhibition of enzymes that destroy
acetylcholine. These modulating factors may be exerted through neurones
where the mediator is non-cholinergic; these are referred to as indirect effects.
15 Some attempts at treatment have focussed on the role of other mediators such as
5-hydroxytryptamine, which is a mediator in other areas of brain, such as the
mid-brain nuclei. However, since fibres from these areas are projected forward
into the cerebral cortex where the primary transmitter is acetylcholine, attention
has focussed on the management of this mediator in the search for appropriate
20 therapeutic agents.

Cholinergic strategies for the treatment of AD/SDAT have been directed
at several points along the pathway of formation, synaptic release and removal
of released acetylcholine.

25 One approach involves treatment with high doses of lecithin and other
precursors of acetylcholine. This is of limited use in producing sustained
improvements in cognitive performance.

30 Another approach involves the use of vegetable drugs such as Polygalae
root extract, which has been shown to enhance choline-acetylcholine transferase

(CAT) activity and nerve growth factor (NGF) secretion in brain. Oral administration of NGF has no effect on central nervous system neurons because it is a high molecular weight protein that cannot pass through the blood-brain barrier. However, agents which can pass through the blood-brain barrier and have a stimulating effect on NGF synthesis in the central nervous system have been proposed for the improvement of memory-related behaviour.

The results of a third clinical approach, which uses cholinesterase inhibitors such as tacrine hydrochloride, have been marginally more positive than the above. Substances obtained from plants used in Chinese and Western medicine, for example huperzine, galanthamine, and physostigmine have all been shown to be of some - although limited - benefit in the treatment of AD/SDAT in clinical studies and also in laboratory models. All of these substances are inhibitors of acetylcholine esterase (AChE). In patients with AD/SDAT, there may be reduced synthesis of acetylcholine (ACh), reduced efficiency in release of ACh from presynaptic stores, and a decrease in the number or function of postsynaptic (M_1) receptors. Reductions in pre-synaptic M_2 receptors have also been shown. The beneficial effect of AChE inhibitors is attributed to enhancement of acetylcholine levels at synapses in brain by slowing down the destruction of released transmitter.

Compositions which modulate cholinergic function are known to affect memory and recall. For example, nicotine stimulates nicotinic acetylcholine receptors, and the short lived memory enhancing effects of cigarette smoking are thought to be due to the effect of nicotine. Scopolamine, an antagonist of acetylcholine, will produce amnesia and impaired cognitive function manifesting in psychomotor tests as a prolongation of simple reaction times, possibly as a result of impaired attention, and is used for this purpose as an adjunctive analgesic treatment. The amnesic effect of scopolamine can be antagonised by nicotine.

There are two families of nicotinic receptor subtypes (α and β), and each includes four subgroups which differ in ligand specificity. The role of nicotinic receptors in the CNS is not well understood at the molecular level. It is possible that agents binding to nicotinic receptors may modify the rate of turnover at muscarinic receptor sites in brain. Nicotinic receptors are ligand-gated ion channels, and their activation causes a rapid (millisecond) increase in cellular permeability to Na^+ and Ca^{++} , depolarisation and excitation.

Another class of cholinergic receptors can be stimulated by muscarine. Such muscarinic (M) receptors are G protein-coupled receptors. Responses of muscarinic receptors are slower; they may be excitatory or inhibitory. They are not necessarily linked to changes in ion permeability. Five types of muscarinic receptors have been detected by cholinergic receptor cloning, and are designated as m_1 - m_5 . Pharmacological effects are associated with four of the cloned receptors and they are designated as M_1 - M_4 based on pharmacological specificity.

Using specific receptor proteins and monoclonal antibodies, it has been possible to further localise muscarinic receptors in brain as m_1 (postsynaptic) and m_2 (presynaptic). In heart, M_2 receptors are postsynaptic. Presynaptic muscarinic receptors are thought to be inhibitory, the binding of ACh to these receptors attenuating the release of further ACh to provide a negative feedback mechanism for ACh release. Selective M_2 receptor antagonists which are preferentially distributed to the brain may therefore be useful in treating Alzheimer's disease.

It is known that, in disease states such as AD/SDAT, there is general neuronal loss and deficits in cholinergic nerve function. It has been speculated that the high affinity nicotinic binding sites in the remaining cholinergic neurons might be converted to low affinity binding sites in treating such diseases, thereby sustaining transmitter release. By lowering the affinity of the nicotinic

binding sites; a quick desensitising process is avoided.

Agonist activation at nicotinic receptors in brain has rapid onset and offset. A decreased affinity of the nicotinic receptors will reduce the desensitisation process. Schwarz R.D. et al (J. Neuro Chem 42, (1984), 1495-8) have shown that nicotine binding sites are presynaptically located on cholinergic (and also 5-hydroxytryptaminergic and catecholaminergic) axon terminals. A change in high affinity binding sites on AD/SDAT may also induce a change in the modulatory effect the nicotinic binding sites may have on other transmitter systems.

Presynaptic cholinergic mechanisms are also under inhibitory control by GABAergic neurons and this inhibition is thought to be intensified in AD/SDAT. Removal or reduction of this inhibition intensifies presynaptic cortical cholinergic activity and enhances cognitive processing.

The interactions of interneuronal fibres innervated by nicotine (reducing binding affinity), and dis-inhibition of GABAergic fibres both have a presynaptic locus.

This is a simplistic model of central transmission, but provides a framework for understanding the attempts which have been made to increase the effective concentration of acetylcholine in central synapses. This further illustrates the concept of direct and indirect action. There are disadvantages attaching to the three conventional therapeutic approaches to AD/SDAT treatment mentioned above: ACh precursor supplementation, agonist replacement and acetylcholine esterase inhibition. These treatments may result in a short-term increase in the availability of ACh which may activate feedback mechanisms resulting in the desensitisation of postsynaptic receptors. On theoretical grounds, long term benefits would not be predicted and when treatment is interrupted, any benefits in management of AD/SDAT and AAMI

disappear and the condition may even be aggravated.

It has been shown that a compound with M_1 agonist and M_2/M_3 antagonist activity improved cognitive performance in SDAT patients (Sramak et al, Life Sciences vol. 2, No. 3, 195-202, 1997). However, this compound causes unacceptable cholinergic side effects, such as fatigue, diarrhoea and nausea.


A more radical approach to AD/SDAT and AAMI aims to increase the number of postsynaptic (M_1) receptors, in brain. It is known from Chinese Patent No. CN1096031A, that sarsasapogenin (SaG) can up-regulate M_1 cholinergic receptors and also down-regulate (i.e. move towards normal levels of) β -adrenergic receptors, the number of which may be pathologically-raised in AD/SDAT.

Patent applications have been published which claim the usefulness of a number of steroid sapogenins having spirostane, furo-spirostane, spirosolane or solanidine structures in the treatment of diseases including SDAT. Two patent publications are of particular relevance here: Chinese patent publication No CN1096031A claims the use of the spirostane sapogenin, sarsasapogenin, in the treatment of SDAT. The disclosure in this document, however, is brief. The other document of relevance is patent publication DE 4303214A1 which claims the use of a very wide range of saponins and sapogenins in the treatment of a whole range of diseases that the inventors consider to be of viral origin. This disclosure is however of dubious value in that it is well recognised that there is no infective element to a very large number of the conditions that are characterised by deficient synaptic transmission and thus the basic premise of the alleged invention is flawed. In addition they present no data of any kind that allows one skilled in the art to be able select a preferred compound from the large number that are claimed.

The inventors have found that smilagenin (SMI) and anzurogenin D (AZD) exhibit the ability to regulate receptors. In particular, these compounds - and especially SMI - have been found to increase the number of M2 receptors in the brain. Thus, according to one aspect of the invention, there is provided the use of smilagenin and/or anzurogenin D in the manufacture of a medicament for the treatment of a condition characterised by a deficiency in postsynaptic membrane-bound receptor number or function.

Those skilled in the art will be aware of the relationship between saponins and their sapogenins, and that the desired effects of sapogenins can be exhibited in patients by administration of the corresponding saponins, or a mixture thereof. Hydrolysis of at least a proportion of saponin occurs in the gastrointestinal tract. The skilled man will also be aware of the epimerisation of certain sapogenins under conditions of acid hydrolysis.

The sapogenins of interest in this invention are of the following general formula:



With reference to this general formula, SMI and AZD have the structure indicated in the Table below:

Compound	A/B ring Cis/Trans/ unsaturation	C25 methyl stereochemistry (R or S)	Hydroxyl group(s) on Spirostane ring
Smilagenin	Cis	R	3 β -OH
Anzurogenin-D	Trans	R	3 β -OH, 5 α -OH, 6 β -OH

5

The saponins and sapogenins of interest in the present invention occur naturally in a range of plant species, notably from the genera *Smilax*, *Asparagus*, *Anemarrhena*, *Yucca* and *Agave*. The species presently of greatest interest include *Smilax regelii* Kilip & Morton - commonly known as Honduran sarsaparilla; *Smilax aristolochiaefolia* Miller - commonly known as Mexican sarsaparilla; *Smilax ornata* Hooker - commonly known as Jamaican sarsaparilla; *Smilax aspera* - commonly known as Spanish sarsaparilla; *Smilax glabra* Roxburgh; *Smilax febrifuga* - Kunth - commonly known as Ecuadorian or Peruvian sarsaparilla; *Anemarrhena asphodeloides* Bunge; *Yucca schidigera* Roez l ex Ortgies; and *Yucca brevifolia* Engelm.

10
15

According to a further aspect of the present invention, there is provided a pharmaceutical composition having cognitive function enhancing properties which comprises an effective amount of smilagenin and/or anzurogenin D.

20

In another aspect, the invention provides a pharmaceutical composition having cognitive function enhancing properties which comprises an effective amount of smilagenin and/or anzurogenin D in the form of an extract derived from a plant of the genus *Smilax*, *Asparagus*, *Anemarrhena*, *Yucca* or *Agave*.

25

It will be appreciated that the invention embraces within its scope the use

of the compositions defined above. Thus, according to a fifth aspect, the present invention provides a method of enhancing cognitive function which comprises administering to a human or animal an effective dosage of a composition of the invention.

5

The invention also provides a method of enhancing cognitive function in a human or non-human animal, which comprises administering an effective dose of smilagenin and/or anzurogenin D.

10

As used herein, the term "cognitive function" refers to functions such as thinking, reasoning, remembering, imagining and learning.

15

In identifying compounds that would have use in the treatment of SDAT and other diseases characterised by reductions in receptor numbers or synaptic transmission, the inventors have given consideration to the need to identify compounds that would have the desired effect but would be devoid of any oestrogenic effects, as these would be unacceptable, particularly in male patients. A number of the compounds claimed to have activity in patent application DE 4303214A1 have marked oestrogenic activity and are therefore unacceptable. Smilagenin and anzurigenin D, however, do not display oestrogenic activity. In addition these two compounds were tested at other steroid receptors and neither compound was found to have any activity at any of the following receptors:

20

Progesterone
Glucocorticoid
Testosterone

25

The selected compounds have also been tested for their activity in a number of in-vitro assays. The assays/experiments that were considered of key importance in determining possible activity in the elevation of membrane bound

receptor numbers were as follows:

1. Chinese hamster ovary (CHO) cells transfected with the a DNA fragment coding for a muscarinic receptor. The cell line used for the majority of the experiments was a cell line expressing the m2 receptor.
2. The effects of muscarinic receptor expression in cultured cell lines of neuronal origin were investigated.
3. Cultured cardiac muscle cells obtained from neonatal Sprague Dawley rats. The cardiac muscle cells express muscarinic receptors, typically m2. The level of these receptors falls on prolonged culture and the effects of compounds of interest in preventing the fall in receptor numbers was investigated.

The methods and the results of these experiments are now described in turn.

1 CHO cell line experiments

The effects of various compounds on the expression of m2 receptors on CHO cells transfected with DNA for the m2 receptor were investigated. Receptor numbers were assayed using tritiated QNB binding and subtracting non-specific binding. Compounds were dissolved in DMSO and DMSO was used as a control. Compounds were tested at a range of final concentrations. Compounds were also tested in the presence and absence of tamoxifen to try to distinguish an oestrogen receptor mediated mechanism. The results are summarised in the Table 2 below, where the compounds used in the invention appear in bold, and data on other sapogenins is given for comparative purposes:

Table 2 Effects of compounds on the expression of m_2 receptors on CHO cells

Compound	Molar concentration of compound	Effect on receptor expression – given as % increase compared to control (negative values in brackets)
Sarsasapogenin	10^{-5}	34
	10^{-6}	(14)
Anzurogenin D	10^{-5}	22
	10^{-6}	(26)
Sisalgenin	10^{-5}	NS
	10^{-6}	NS
Smilagenin	10^{-5}	57
	10^{-6}	18
Diosgenin	10^{-5}	NS
	10^{-6}	NS
Ruscogenin	10^{-5}	(22)
	10^{-6}	NS
Tigogenin	10^{-5}	NS
	10^{-6}	NS

20

NS = No significant effect

Thus the experiments indicate that smilagenin and anzurogenin D were able to increase the number of muscarinic receptors expressed on the surface of CHO cells cultured in-vitro. The effect was not antagonised by tamoxifen, indicating that the mechanism involved did not involve the oestrogen receptor.

5

2 Effects of compounds on cell survival

10

15

20

Other in vitro assays have been employed to establish the effects of smilagenin and anzurogenin D. In particular various neuroblastoma cell lines including SKN-SN and SH-SY5Y cells as well as phaeochromocytoma cell lines have been cultured in vitro in the presence of β -amyloid fragments or serum depletion. A number of techniques to demonstrate the effectiveness of the compounds in protecting the cultured cells were investigated. These techniques included Trypan blue exclusion, chemiluminescence and release of lactate dehydrogenase. Of most interest was the observation that incubation of cells, in particular PC12 cells, with β -amyloid reduced the number of muscarinic receptors measured using radio-labelled ligand binding techniques. This reduction in receptor numbers was found to be ameliorated by smilagenin and by anzurogenin D.

3 Effects of compounds on cultured cardiac muscle cells.

25

30

Cardiac muscle cells were isolated from the ventricular muscle of neonatal Sprague Dawley rats using standard techniques. Cells were cultured in vitro and muscarinic receptor numbers expressed on cell surfaces membrane fragments after homogenisation of cells harvested at various time points were estimated using specific binding of tritiated QNB. Preliminary experiments demonstrated that the number of receptors expressed tended to decline after 10 days of culture. The experiments were therefore designed to investigate the effects of the various compounds in inhibiting this decline in receptor numbers.

The results of these experiments are summarised in Table 3, where the compounds used in the invention appear in bold, and data on other sapogenins is given for comparative purposes:

5 **Table 3: Effects of various compounds on muscarinic receptor expression on cultured cardiac muscle cells**

Compound	Concentration of compound causing a significant increase in number of receptors expressed on neonatal cardiac muscle after 10 days in vitro culture
Diosgenin	NS
10 Anzurogenin D	$10^{-6}M$
Ruscogenin	NS
Sarsasapogenin	$10^{-5}M$
Tigogenin	NS
Astragaloside	$10^{-5}M$
15 Smilagenin	$10^{-6}M$

NS = No significant effect

20 It is speculated here that the effect of the active compounds claimed in this patent may operate through an effect on G protein and that the effects on receptor numbers are secondary to an effect on G-protein. When a membrane bound G-protein linked receptor is stimulated two basic sets of events are initiated: the effector response; and the internalisation of the receptor. The

-14-

subsequent processing of the receptor to the state where it is again in a form on the cell surface or other membrane surface where it can interact with another receptor ligand appears to be subject to a number of factors. A number of these factors or mechanisms appear to be G-protein linked. There is evidence that activation of m_3 receptors may have an effect on G-protein expression or levels. It is speculated that the actions of the compounds described in this patent may due to an interaction in the processes of receptor regeneration, G-protein linkage or G-protein homeostasis.

An alternative hypothesis is that the compounds are increasing the synthesis or release or a decreased rate of degradation of neurotropic factors such as brain derived growth factor and/or nerve growth factor. These effects on growth factors might be due to an effect of the compound on a cytosolic or nuclear receptor or the binding of a compound to a promoter region with a consequent effect directly on the rate of production of mRNA for the growth factor or as a consequence of increasing the production of another material factor such as G-protein or finally the effects may be secondary to an effect on receptor or G-protein procession.

CLAIMS:

1. The use of smilagenin in the manufacture of a medicament for the treatment of a condition characterised by a deficiency in postsynaptic
5 membrane-bound receptor number or function.

2. The use of anzurogenin D in the manufacture of a medicament for the treatment of a condition characterised by a deficiency in postsynaptic
10 membrane-bound receptor number or function.

3. A pharmaceutical composition having cognitive function enhancing properties which comprises a pharmacologically effective amount of smilagenin and/or anzurogenin D.

4. A pharmaceutical composition having cognitive function enhancing properties which comprises a pharmacologically effective amount of smilagenin and/or anzurogenin D in the form of an extract derived from a plant of the
15 genus Smilax, Asparagus, Anemarrhena, Yucca or Agave.

5. A method of enhancing cognitive function which comprises administering to a human or animal an effective dosage of a composition as
20 claimed in claim 3 or 4.

6. A method of enhancing cognitive function in a human or non-human
25 animal, which comprises administering an effective dose of smilagenin and/or anzurogenin D.

7. A method of enhancing cognitive function in a patient suffering from age-related cognitive disfunction, which comprises administering to the patient a
30 pharmacologically effective dose of smilagenin and/or anzurogenin D.

8. A method as claimed in claim 5 or 6, which is for the treatment of Alzheimer's disease or a senile dementia of the Alzheimer's type.

5 9. A method as claimed in claim 5, 6, 7 or 8, substantially as hereinbefore described.

10. A composition for use in a method as claimed in any one of claims 5 to 9, substantially as hereinbefore described.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.